





# AP Bio

# Unit Reviews

Ecology

@apbiopenguins



**AP Biology students are  
penguins because they are  
Dressed for Success!**

**You are now an AP Bio Penguin!**



## Resource Reminders:

Daily Review on IG stories

374 page Review Guide on Weebly

Recorded FRQ Fridays on YouTube

120+ Quizizz Games on Weebly

Review PowerPoints on Weebly

Weebly: [www.apbiopenguins.weebly.com](http://www.apbiopenguins.weebly.com)



# Today's Plan

Energy Flow

Population Ecology

Community Ecology

Practice Questions

Unit 8 Q&A





# Body Temperature

## Endotherm

Maintains body temperature through metabolism

## Ectotherm

Maintains body temperature through behaviors (bask in sunlight, aggregation)



# Energy Flow

Organisms use energy to maintain, organize, grow and reproduce


## Autotroph

Capture energy from physical or chemical source

- Photosynthetic - sunlight
- Chemosynthetic

small inorganic molecules in environment (sometimes w/o oxygen)

## Trophic Structure



## Heterotroph

Capture energy present in carbon compounds produced by other organisms

Metabolize carbohydrates, lipids, and proteins (notice: not nucleic acids) for energy through hydrolysis

## Changes in Availability

### Change in Energy Resource

Affects number and size of trophic levels

### Change in Producer Level

Affects number and size of trophic levels

# Animal Behavior

## Communication

Signaling allows for changes in behaviors of organisms to allow for differential reproductive success

### Types of Communication:

- Visual
- Auditory
- Electrical
- Chemical


### Function:

- Indicate Dominance
- Foraging (Finding Food)
- Establish Territory
- Ensure Reproductive Success


## Altruistic Behaviors

Reduces individual fitness but increases inclusive fitness.

## Examples of Animal Communication



Fireflies glow to attract mates.



Peacocks use their elaborate tails during courting rituals.



Cobras inflate their hood to scare other creatures.

**Visual**



Elephants use their trunks to talk to other herds over long distances.



Male whales use their song to communicate with females.



Wolves howl to call to other wolves in the pack.

**Auditory**



Dogs lick their pups to bond, clean and stimulate their development.



Baboons use touch to show affection and groom each other.



Horses kick other horses to establish dominance.

**Tactile**



Cats rub against objects to mark them with their scent.



Ants use pheromone trails to follow each other.



Skunks use their signature smell to deter predators.

**Chemical**

# Animal Behavior

## Communication

Signaling allows for changes in behaviors of organisms to allow for differential reproductive success

### Types of Communication:


- Visual
- Auditory
- Electrical
- Chemical

### Function:

- Indicate Dominance
- Foraging (Finding Food)
- Establish Territory
- Ensure Reproductive Success

## Altruistic Behaviors

Reduces individual fitness but increases inclusive fitness.



## Intersexual Selection

Reproductive behaviors to attract a mate  
Individuals of one sex choose members of the opposite sex

### Examples

- Blue Footed Booby – mating dance (visual)
  - Frogs – croaking (auditory)
  - Pheromones – (chemical)

## Intrasexual Selection

Reproductive behaviors to indicate dominance and compete for access to mates

### Examples

- Deer: antler size
- Horned Beetles: strength and size of “horn”

# Population Ecology

## Exponential Growth

Unlimited growth of population

$$r = b - d$$

rate of increase = birth rate - death rate

$$\frac{dN}{dt} = rN$$

### Example:


If a population has **400** individuals with a rate of increase of **0.5**, how many individuals after **2** generations?

$$\frac{dN}{dt} = (0.5)(400) = 200$$

After generation **1**: **600**

$$\frac{dN}{dt} = (0.5)(600) = 300$$

After generation **2**: **900**



### Density Dependent Factors

Factors that intensifies as population increases

Ex: competition, predation, disease

### Density Independent Factors

Factors that affect all individuals regardless of size, population, density

Ex: natural disasters, human activity

## Logistic Growth

Population size limited by carrying capacity

$$\frac{dN}{dt} = rN\left(\frac{K - N}{K}\right) = 200$$

### Example:

If a population has **400** individuals with a rate of increase of **0.5** and a carrying capacity of **800**, how many individuals after **2** generations?

$$\frac{dN}{dt} = (0.5)(400)\left(\frac{800 - 400}{800}\right) = 100$$

After generation **1**: **500**

$$\frac{dN}{dt} = (0.5)(500)\left(\frac{800 - 500}{800}\right) = 75$$

After generation **2**: **575**



# Species Diversity

Simpson's Index: measures biodiversity (species composition and diversity)

$$Simpson\ Diversity = 1 - \sum \left(\frac{n}{N}\right)^2$$

n = total number of organisms of particular species  
N = total of organisms of all species

Species	Number
Sloth	18
Penguin	13
Total	31

$$1 - \left( \left(\frac{18}{31}\right)^2 + \left(\frac{13}{31}\right)^2 \right)$$

$$1 - ((0.58)^2 + (0.42)^2)$$

$$1 - (0.34 + 0.18)$$

$$1 - 0.52 = 0.48$$

# Keystone Species

Organism with disproportionate to their abundance effect, and when they are removed from the ecosystem, the ecosystem often collapses.

Examples: Sea Otter

# Community Ecology




## Interactions

- Predator/Prey (+/-)
- Herbivory (+/-)
- Competition (-/-)

## Symbiosis

- Parasitism (+/-)
- Mutualism (+/+)
- Commensalism (+/0)

## Ecological Relationships



@AmoebaSisters





Removal of the **keystone sea otter** : sea urchins overgraze kelp and destroy the kelp forest community.

## Keystone Species

with a disproportionate impact on their abundance and diversity. When they are removed from the ecosystem, the ecosystem often collapses.

Examples: Sea Otter



# Species Diversity

Simpson's Index: measures biodiversity  
(species composition and diversity)

$$Simpson\ Diversity = 1 - \sum \left(\frac{n}{N}\right)^2$$

n = total number of organisms of particular species  
N = total of organisms of all species

Species	Number
Sloth	18
Penguin	13
Total	31

$$1 - \left( \left(\frac{18}{31}\right)^2 + \left(\frac{13}{31}\right)^2 \right)$$

$$1 - ((0.58)^2 + (0.42)^2)$$

$$1 - (0.34 + 0.18)$$

$$1 - 0.52 = 0.48$$

# Keystone Species

Organism with disproportionate to their abundance effect, and when they are removed from the ecosystem, the ecosystem often collapses.




Examples: Sea Otter

# Community Ecology


## Interactions

- Predator/Prey (+/-)
- Herbivory (+/-)
- Competition (-/-)
- Symbiosis
  - Parasitism (+/-)
  - Mutualism (+/+)
  - Commensalism (+/0)

## Ecological Relationships



@AmoebaSisters



## Invasive Species


Organism that is not indigenous, or native, to a particular area with no natural predators and unlimited resources

Examples:

- Zebra Mussel: clogging water way
- Lionfish: venomous species

## Multiple Choice Practice:

Testosterone oxido-reductase is a liver enzyme that regulates testosterone levels in alligators. One study compared testosterone oxido-reductase activity between male and female alligators from Lake Woodruff, a relatively pristine environment, and from Lake Apopka, an area that has suffered severe contamination. The graph above depicts the findings of that study.




The data in the graph best support which of the following claims?

- Environmental contamination elevates total testosterone oxido-reductase activity in females.
- Environmental contamination reduces total testosterone oxido-reductase activity in females.
- Environmental contamination elevates total testosterone oxido-reductase activity in males.
- Environmental contamination reduces total testosterone oxido-reductase activity in males.



Testosterone oxido-reductase activity was compared between males and females at two sites: a relatively pristine environment (Woodruff) and a contaminated environment (Apopka). The graph above depicts the results.



alligators. One study from Lake Woodruff, a relatively pristine environment, and a study from Apopka, a site of severe contamination. The graph above depicts the results.


The data in the graph best support which of the following claims?

- Environmental contamination elevates total testosterone oxido-reductase activity in females.
- Environmental contamination reduces total testosterone oxido-reductase activity in females.
- Environmental contamination elevates total testosterone oxido-reductase activity in males.
- Environmental contamination reduces total testosterone oxido-reductase activity in males.



## Multiple Choice Practice:

Testosterone oxido-reductase is a liver enzyme that regulates testosterone levels in alligators. One study compared testosterone oxido-reductase activity between male and female alligators from Lake Woodruff, a relatively pristine environment, and from Lake Apopka, an area that has suffered severe contamination. The graph above depicts the findings of that study.



The data in the graph best support which of the following claims?

- a. Environmental contamination elevates total testosterone oxido-reductase activity in females.
- b. Environmental contamination reduces total testosterone oxido-reductase activity in females.
- c. Environmental contamination elevates total testosterone oxido-reductase activity in males.
- d. Environmental contamination reduces total testosterone oxido-reductase activity in males.




## Multiple Choice Practice:

Beaked whales feed at various depths, but they defecate at the ocean's surface.

Nitrogen-rich whale feces deposited in surface waters supply nutrients for algae

that are eaten by surface dwelling fish. Which of the following best predicts what would happen if the whale population decreased?

- a. There would be a reduction in surface nitrogen concentration, which would cause an algal bloom.
- b. The surface fish populations would decline due to reduced populations of algae.
- c. The remaining whales would accumulate mutations at a faster rate.
- d. The remaining whales would be forced to forage in the deepest parts of the ocean.



## Multiple Choice Practice:

Scientists have found that the existing populations of a certain species of amphibian are small in number, lacking in genetic diversity, and separated from each other by wide areas of dry land. Which of the following human actions is most likely to improve the long-term survival of the amphibians?


- a. Cloning the largest individuals to counteract the effects of aggressive predation
- b. Reducing the population size by one-fifth to decrease competition for limited resources
- c. Constructing a dam and irrigation system to control flooding
- d. Building ponds in the areas of dry land to promote interbreeding between the separated populations





## Free Response Practice (2013 #1c/d):

(c) The experiment described above is repeated with ripe bananas at one end and unripe bananas at the other end. Once again the positions of the flies are observed and recorded every minute for **10** minutes. The positions of flies after **1** minute and after **10** minutes are shown in the table below.



DISTRIBUTION OF FLIES IN CHOICE CHAMBER

Time (minutes)	Position in Chamber		
	End with Ripe Banana	Middle	End with Unripe Banana
1	21	18	21
10	45	3	12

Perform a chi-square test on the data for the **10**-minute time point in the banana experiment. Specify the null hypothesis that you are testing and enter the values from your calculations in the table below.

(d) Explain whether your hypothesis is supported by the chi-square test and justify your explanation.



# Free Response Practice (2013 #1c/d):

Perform a chi-square test on the data for the 10-minute time point in the banana experiment. Specify the null hypothesis that you are testing and enter the values from your calculations in the table below.

DISTRIBUTION OF FLIES IN CHOICE CHAMBER

Time (minutes)	Position in Chamber		
	End with Ripe Banana	Middle	End with Unripe Banana
1	21	18	21
10	45	3	12

Null Hypothesis:

	Observed (o)	Expected (e)	$(o-e)^2/e$
End with ripe banana			
Middle			
End with unripe banana			
Total			



# Free Response Practice (2013 #1c/d):

(d) Explain whether your hypothesis is supported by the chi-square test and justify your explanation.

## Null Hypothesis:

The presence of the banana (ripe or unripe) has no effect on the location of the flies in the choice chamber. The flies will be equally distributed between the regions of the choice chamber.

	Observed (o)	Expected (e)	$(o-e)^2/e$
End with ripe banana	45	20	31.25
Middle	3	20	14.45
End with unripe banana	12	20	3.2
Total	60	60	48.9

## Chi-Square Table

p value	Degrees of Freedom							
	1	2	3	4	5	6	7	8
0.05	3.84	5.99	7.81	9.49	11.07	12.59	14.07	15.51
0.01	6.63	9.21	11.34	13.28	15.09	16.81	18.48	20.09

$$48.9 > 5.99$$

At a  $p = 0.05$  with 2 degrees of freedom, the calculated value is greater than the table "critical" value so we will reject the null hypothesis. The banana ripeness has an affect on the fly distribution.



## Free Response Practice (2017 #4):

The table below shows how much each organism in an aquatic ecosystem relies on various food sources. The rows represent the organisms in the ecosystem, and the columns represent the food source. The percentages indicate the proportional dietary composition of each organism. High percentages indicate strong dependence of an organism on a food source.

DIETARY COMPOSITION OF ORGANISMS IN AN AQUATIC ECOSYSTEM

Organism	Food Source (% of diet)				
	Algae	Stoneflies	Midges	Hellgrammites	Caddisflies
Algae					
Stoneflies			90		10
Midges	100				
Hellgrammites		20	10		70
Caddisflies	70		30		




# Free Response Practice (2017 #4):

(a) Based on the food sources indicated in the data table, construct a food web in the template below. Write the organism names on the appropriate lines AND draw the arrows necessary to indicate the energy flow between organisms in the ecosystem.

DIETARY COMPOSITION OF ORGANISMS IN AN AQUATIC ECOSYSTEM

Organism	Food Source (% of diet)				
	Algae	Stoneflies	Midges	Hellgrammites	Caddisflies
Algae					
Stoneflies			90		10
Midges	100				
Hellgrammites		20	10		70
Caddisflies	70		30		



# Free Response Practice (2017 #4):

(b) In an effort to control the number of midges, an area within the ecosystem was sprayed with the fungus *Metarhizium anisopliae*, which significantly decreased the midge population. Based on the data in the table, predict whether the spraying of fungus will have the greatest short-term impact on the population of the stoneflies, the caddisflies, or the hellgrammites. Justify your prediction.

DIETARY COMPOSITION OF ORGANISMS IN AN AQUATIC ECOSYSTEM

Organism	Food Source (% of diet)				
	Algae	Stoneflies	Midges	Hellgrammites	Caddisflies
Algae					
Stoneflies			90		10
Midges	100				
Hellgrammites		20	10		70
Caddisflies	70		30		

## Prediction (1 point)

- Stoneflies

## Justification (1 point)

- Stoneflies have a higher dependence on the midges than do the hellgrammites and caddisflies.
- Midges are 90 percent of the stonefly diet, while 30 percent of the caddisfly and 10 percent of the hellgrammite diet are midges.



# Free Response Practice (2018 #5):

Some birds, including great spotted cuckoos, lay their eggs in the nests of other birds, such as reed warblers. The warbler parents raise the unrelated chicks and provide them with food that would otherwise be given to their biological offspring. A researcher conducted an investigation to determine the type of relationship between warblers and cuckoos in an environment without predators. The researcher found that nests containing only warblers were more likely to be successful than nests containing warblers and cuckoos (data not shown). A successful nest is defined as a nest where at least one chick becomes an adult warbler.

In some geographic areas, several species of nest predators are present. Researchers have found that cuckoo chicks, while in the nest, produce a smelly substance that deters nest predators. The substance does not remain in the nest if cuckoo chicks are removed. Figure 1 shows the probability that nest containing only warblers or containing both warblers and cuckoos will be successful in an environment with predators. In a follow-up experiment, the researchers added cuckoos to a nest that contained only warblers (group 1) and removed from a nest containing warblers and cuckoos (group 2).




Figure 1. Probability of nest success in an environment with predators

# Free Response Practice (2018 #5):

(a) Describe the symbiotic relationship that exists between the cuckoo and warbler in an environment without predators.

Cuckoo are parasites to the warblers because the cuckoo is benefited while the warbler is harmed in the relationship.

Recall: Cuckoos lay their eggs in the nests of warblers. The warbler parents raise the unrelated chicks and provide them with food that would otherwise be given to their biological offspring.





# Free Response Practice (2018 #5):

(b) On the template provided, draw bars in the appropriate locations to predict the relative probability of success for the nest in the presence of predators where:

- \* the cuckoos were added to the nest containing only warblers (group 1)
- \* the cuckoos were removed from the nest containing warblers and cuckoos (group 2)



# Free Response Practice (2018 #5):

(c) Identify the symbiotic relationship that exists between the cuckoo and the warbler in the presence of predators.

Cuckoo bird and the warbler are in a mutualistic relationship.

Recall: Researchers have found that cuckoo chicks, while in the nest, produce a smelly substance that deters nest predators. The substance does not remain in the nest if cuckoo chicks are removed.



Q & A



## CRAM Sessions

**Saturday 5/11 @3pm**

**AP Bio Penguins**

**Wednesday 5/15 @8pm**

**MarcoLearning**



## Check Out These Additional Resources

**MarcoLearning**  
**APsoluteRecAP**  
**Etch.com**





**@apbiopenguins**



**@apbiopenguins**



**AP Bio Insta-Review**

**Follow us on your favorite  
social media channels!**

